

2/29/04

12/18/04

L21 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2003:868614 CAPLUS  
 DN 139:371877  
 TI Positive-working resist composition  
 IN Sasaki, Tomoya; Mizutani, Kazuyoshi; Kanna, Shinichi  
 PA Fuji Photo Film Co., Ltd., Japan  
 SO Jpn. Kokai Tokkyo Koho, 60 pp.  
 CODEN: JKXXAF

DT Patent  
 LA Japanese

FAN. CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2003316007	A2	20031106	JP 2002-126433	20020426
	US 2003232277	A1	20031218	US 2003-422789	20030425
PRAI	JP 2002-126433	A	20020426		
	JP 2002-223234	A	20020731		
	JP 2002-223386	A	20020731		

AB The pos.-working resist composition comprises a resin which increases its solubility

in an alkali developer upon reaction with an acid and has  $\geq 1$  repeating unit selected from I and II (R1,2 = H, halo, cyano, etc.; L1 = divalent bonding group; m1, m2 = 0, 1; R3,4 = alkyl, aryl, etc.; Rb = H, organic group, halo; l = 0-3; and L2 = single bond, divalent bonding group) and  $\geq 1$  repeating unit such as III (R11-16 = H, alkyl, etc.), a compound generating an acid upon receiving an active ray or a radiation. The use of the resin exhibited high optical transparency at  $\leq 160$  nm.

IT 622840-85-1 622840-98-6

RL: TEM (Technical or engineered material use); USES (Uses)  
 (resin in pos.-working resist composition)



RN 622840-85-1 CAPLUS

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 2,2,2-trifluoro-1-[4-[2,2,2-trifluoro-1-(methoxymethoxy)-1-(trifluoromethyl)ethyl]cyclohexyl]-1-(trifluoromethyl)ethyl ester, polymer with 1-methylethyl 4-ethenylbenzenesulfonate (9CI) (CA INDEX NAME)



CM 1

CRN 610301-02-5  
 CMF C18 H17 F15 O4

Sec. 143, Cal. 13





Both have ART above see 143 pg 13




CM 2

CRN 79293-23-5  
 CMF C11 H14 O3 S

my case sec. 60 pg 9  
 A + CO<sub>2</sub>






RN 622840-98-6 CAPLUS

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1-[4-[1-[(1,1-dimethylethoxy)methoxy]-2,2,2-trifluoro-1-(trifluoromethyl)ethyl]cyclohexyl]-2,2,2-trifluoro-1-(trifluoromethyl)ethyl ester, polymer with methyl 4-ethenylbenzenesulfonate and 2,2,2-trifluoro-1-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]cyclohexyl]-1-(trifluoromethyl)ethyl 2-(trifluoromethyl)-2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 622840-97-5


CMF C21 H23 F15 O4



CM 2

CRN 479072-83-8


CMF C16 H13 F15 O3



CM 3

CRN 16736-97-3

CMF C9 H10 O3 S



L21 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2005 ACS on STN  
 AN 2001:372174 CAPLUS  
 DN 134:368010  
 TI UV-absorbing polymers bearing 2-hydroxybenzophenone or  
 2'-hydroxyphenylbenzotriazole residues  
 IN Aoyama, Masato  
 PA Mitsubishi Chemical Corp., Japan  
 SO Jpn. Kokai Tokkyo Koho, 6 pp.  
 CODEN: JKXXAF  
 DT Patent  
 LA Japanese  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2001139640	A2	20010522	JP 1999-321992	19991112
PRAI	JP 1999-321992		19991112		

AB The polymers, exhibiting high water solubility and being useful for UV stabilizers especially in waterborne emulsion coatings, comprise (i) UV-absorbing vinylic monomers bearing 2-hydroxybenzophenone or 2'-hydroxyphenylbenzotriazole residues and (ii) water-soluble vinylic monomers bearing anion-dissociative groups. The monomer (ii) may be 2-acrylamidopropanesulfonic acid, 2-acrylamido-2-methylpropanoic acid, 3-acrylamido-2,4,4-trimethylpentanesulfonic acid, 2-acrylamido-2-(4-tolyl)ethanesulfonic acid, 4-styrenesulfonic acid, and/or their salts. Thus, 1:3 (g) 2-hydroxy-4-(2-methacryloyloxy)ethoxybenzophenone-sodium p-styrenesulfonate copolymer showed water solubility ≥25%, no gelation in blending with Rikabond ES 52 (anionic aqueous emulsion coating), and <380-nm UV absorption.

IT **339994-71-7**  
 RL: POF (Polymer in formulation); TEM (Technical or engineered material use); USES (Uses)  
 (macromol. UV absorbers bearing benzophenone or benzotriazole groups and showing good solubility to anionic aqueous coatings)


RN 339994-71-7 CAPLUS

CN 2-Propenoic acid, 2-methyl-, 3-(2H-benzotriazol-2-yl)-4-hydroxyphenyl ester, polymer with sodium 4-ethenylbenzenesulfonate (9CI) (CA INDEX NAME)

CM 1

CRN 132288-91-6


CMF C16 H13 N3 O3





CM 2

CRN 2695-37-6

CMF C8 H8 O3 S . Na




=>




Text content in the middle section, including various symbols and characters.








b.































𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰  
 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰  
 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰  
 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰 𐌲𐌹𐌿𐌸𐌰

















● ♀ ♂ ♀ ●



Chemical structures 1 through 7 are shown:

- 1: A vertical chain starting with a  $\text{CH}_2$  group, followed by a carbon atom bonded to  $\text{G}_1$  and a carbonyl group ( $\text{C}=\text{O}$ ). This is followed by an oxygen atom, a  $[\text{CH}_2]_{1-20}$  chain, and a cyclohexane ring. The ring is labeled with  $\text{CH}$  at the top and bottom, and  $\text{CH}_2$  on the sides. A  $\text{G}_3$  group is attached to the bottom  $\text{CH}$  of the ring.
- 2: A central carbon atom bonded to two  $\text{CH}_3$  groups and a  $\text{CH}_2$  group.
- 3: A central carbon atom bonded to two  $\text{CH}_3$  groups and a  $\text{CH}_2$  group.
- 4: A central carbon atom bonded to two  $\text{CF}_3$  groups and a  $\text{OH}$  group.
- 5: A central carbon atom bonded to two  $\text{CF}_3$  groups and a  $\text{CH}_2$  group.
- 6: A central carbon atom bonded to two  $\text{CF}_3$  groups and a  $\text{CH}_2$  group.
- 7: A central carbon atom bonded to two  $\text{CF}_3$  groups and a  $\text{CH}_2$  group.

[illegible]





















